



ehponline.org

ENVIRONMENTAL HEALTH PERSPECTIVES

In Utero Pesticide Exposure and Leukemia in Brazilian Children < 2 Years of Age

Jeniffer Dantas Ferreira, Arnaldo César Couto,
Maria do Socorro Pombo-de-Oliveira, Sergio Koifman, and the
Brazilian Collaborative Study Group of Infant Acute Leukemia

<http://dx.doi.org/10.1289/ehp.1103942>

Online 22 October 2012



NIEHS

National Institute of
Environmental Health Sciences

National Institutes of Health
U.S. Department of Health and Human Services

In Utero Pesticide Exposure and Leukemia in Brazilian Children < 2 Years of Age

Jeniffer Dantas Ferreira¹, Arnaldo César Couto¹, Maria do Socorro Pombo-de-Oliveira², Sergio Koifman¹, and the Brazilian Collaborative Study Group of Infant Acute Leukemia³.

¹ Environment and Public Health Post-graduation Program, National School of Public Health, Oswaldo Cruz Foundation (FIOCRUZ), Rio de Janeiro, Brazil

² Pediatric Hematology-Oncology Program, Research Center, Instituto Nacional de Câncer/Rio de Janeiro, Rio de Janeiro, Brazil

Correspondence to:

Sergio Koifman

Environment and Public Health Post-graduation Program, National School of Public Health, Oswaldo Cruz Foundation (FIOCRUZ)

Rua Leopoldo Bulhões 1480, Rio de Janeiro, RJ, 21041-210, Brazil

Telephone number: +55 21 2598-2634

e-mail: koifman@ensp.fiocruz.br;

Brazilian Collaborative Study Group of Infant Acute Leukemia Collaborators

Jozina Maria de Andrade Agareno¹, Flávia Nogueira Serafim Araújo¹, Núbia Mendonça¹, Cynthia Curvello Neves¹, Alejandro Aranciba², Renato Melarangno², Rosania Baseggio³, Atalla Mnayarji³, Reinaldo Del Belo⁴, Jane Dobbin⁴, Silvia Brandalise⁵, Vitória Pinheiro⁵, Lilian M Burlacchini de Carvalho⁶, Eni Guimarães de Carvalho⁶, Tereza Cristina Cardoso⁶, Imaruí Costa⁷, Jose Carlos Cordoba⁸, Isis Q Magalhães⁸, Virginia M Coser⁹, Maria Lucia Lee¹⁰, Flávia Pimenta¹¹, Mara A.D.Pianovsky¹², Terezinha JM Salles¹³, Fernando Werneck¹⁴ and César Bariani¹⁵

¹ Pediatric Hematology-Oncology Service, Hospital Santa Izabel, Salvador, Bahia, Brazil

² Pediatric Hematology-Oncology Service, Hospital Santa Marcelina, Sao Paulo, São Paulo, Brazil

³ Pediatric Hematology-Oncology Service, Hospital Rosa Pedrossian, Campo Grande, Mato Grosso do Sul, Brazil

⁴ Research Center and Hematology Service, Instituto Nacional de Câncer, Rio de Janeiro, Rio de Janeiro, Brazil

⁵ Centro Infantil de Investigações Hematológicas D. Boldrini, Campinas, São Paulo, Brazil

⁶ Pediatric Hematology-Oncology Service, Hospital Martagão Gesteira, Salvador, Bahia, Brazil

⁷ Pediatric Hematology-Oncology Service, Hospital Joana de Gusmão, Florianópolis, Santa Catarina, Brazil

⁸ Hospital de Apoio Brasília, Unidade de Onco-Hematologia Pediátrica, Brasília, Distrito Federal, Brazil

⁹ Departamento de Hematologia, Universidade de Santa Maria, Santa Maria, Rio Grande do Sul, Brazil

¹⁰ Pediatric Oncology Institute- GRAAC, São Paulo, São Paulo, Brazil

¹¹ Hospital Napoleão Laureano, João Pessoa, Paraíba, Brazil

¹² Hospital Pegueno Príncipe, Curitiba, Paraná, Brazil

¹³ Hospital Oswaldo Cruz, CEON, Recife, Pernambuco, Brazil

¹⁴ Pediatric Oncology Section, Hospital dos Servidores do Estado do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

¹⁵ Serviço de Transplantes de Medula do Hospital Araújo Jorge, Goiania, Goiás, Brazil

Running title: Pesticide exposure, Pregnancy and Very Young Leukemia

Keywords: Childhood Cancer, Infant Leukemia, Lactation, Pesticides, Pregnancy.

Acknowledgements: This investigation was supported by the Brazilian National Research Council (CNPq), Instituto Nacional de Câncer-Fundação Ary Frauzino, and the Swiss Bridge Foundation. JDF and ACC have been supported by Oswaldo Cruz Foundation, Ministry of Health of Brazil fellowships. MSPO and SK have been supported by CNPq research grants, #309091/2007-1 and # 577598/2008-2, respectively. The project was granted by INCT- Controle do Cancer; CNPq #573806/2008-0, and the State of Rio de Janeiro Research Foundation (FAPERJ), grant E026/2008.

Conflict of Financial Interests: The authors declare that there are not competing financial conflicts of interests.

ABSTRACT

Background: An association between pesticides exposure and cancer has been suggested. Infant leukemia is a rare neoplasm and its association with maternal pesticide exposure has been poorly explored.

Objectives: To investigate the association between pesticide exposure during pregnancy and leukemia in children < 2 years of age.

Methods: A hospital-based case-control study was carried out in 15 Brazilian hospitals in 1999-2007. Mothers of 252 cases and those of 423 controls were interviewed. Information on pesticide exposures 3 months before pregnancy, throughout pregnancy, and during breastfeeding was obtained. Unconditional logistic regression was used to estimate adjusted odds ratios (adj. ORs) for associations between pesticide exposures and leukemia.

Results: Associations with ever use of pesticides during pregnancy were observed for acute lymphoblastic leukemia, ALL (adj. OR 2.10; 95% CI: 1.14, 3.86) and acute myeloid leukemia, AML (adj. OR 5.01; 95% CI: 1.97, 12.7) in children age 0–11 months, and with ALL (adj. OR 1.88; 95% CI: 1.05, 5.23) at age 12-23 months. According to reported maternal exposure to permethrin, higher risk estimates were verified for children 0-11 months, adj. OR 2.47; 95% CI: 1.17, 5.25 for ALL, and adj. OR 7.28; 95% CI: 2.60, 20.38 for AML. Maternal pesticide exposure related to agricultural activities showed an adj. OR 5.25; 95% CI: 1.83, 15.08 for ALL, and an adj. OR 7.56; 95% CI: 1.83, 31.23 for AML.

Conclusions: These results support the hypothesis that pesticide exposure during pregnancy may be involved in the etiology of acute leukemia in children younger than age 2 years.

INTRODUCTION

Pesticide exposure is a public health concern worldwide. In Brazil, a study conducted in two stages among small-scale fruit farmers revealed that 19% of them had reported at least one poisoning episode (4% in the year previous to interview). Even at this stage, it was concluded that 11% of all episodes were probable cases of acute poisoning according to WHO criteria (Faria et al. 2009). Moreover, pesticide use also has been associated with chronic diseases such as cancer, including childhood leukemia, 0-14 yr. (Bassil et al. 2007; Infante-Rivard and Weichenthal 2007; Turner et al. 2010; Zahm and Ward 1998).

Infant leukemias are rare acute leukemias diagnosed within the first 12 months of age, although some researchers include cases up to 18 months of age (Alexander et al. 2001; Pombo-de-Oliveira and Koifman 2006). DNA alterations during the periconceptual period or pregnancy, which may be caused by chemical substances and other environmental exposures, are potential risk factors for infant leukemias (Buffler et al. 2005; Lafiura et al. 2007). Previous studies have reported associations between childhood leukemia and maternal exposure to pesticides (Ma et al. 2002; Meinert et al. 2000; Menegaux et al. 2006; Rudant et al. 2007; Monge et al. 2007).

However, very few studies of pesticides and leukemia in very young children have been published. Additionally, most studies of association between childhood leukemias and pesticides focused on parents' occupational exposures (Colt and Blair 1998; Monge et al. 2005; Perez-Saldivar et al. 2008; Rudant et al. 2007; Wigle et al. 2009), and a few considered the use of household pesticides during the prenatal period (Lafiura et al. 2007; Meinert et al. 2000; Monge et al. 2007; Zahm and Ward 1998).

A case-control study of risk factors for leukemia in children under 2 years was conducted in Brazil, and an adj. OR 2.18 (95% CI: 1.53, 2.13) was reported in association with maternal exposure to pesticides (Pombo-de-Oliveira and Koifman 2006). The present

investigation aimed to extend such previous exploring analysis on maternal pesticides exposure and leukemia in the offspring.

METHODS

Study Population

This investigation is part of a multicenter study called “Multi-institutional Study of Infant Leukemia: Contribution of Immunomolecular Markers in Distinguishing Different Etiopathogenic Factors”, which focuses on the investigation of biomarkers of leukemia diagnosed in children younger than 2 years in Brazil. Participants (n=675) were recruited from 15 selected hospitals in all geographic areas in the country but the Amazon, including cities in the Southern Region, the Southeast, the Northeast, and the Middle West.

Study Design

This is a hospital-based multicenter case-control study in which controls were frequency matched with leukemia cases according to age (0 – 23 months) and enrolled from the same geographic areas where cases were diagnosed.

Data were obtained by in person interviews carried out between 1999-2007 with mothers of newly diagnosed patients recruited from the Brazilian National Health System centers providing oncologic care for pediatric patients and from general hospitals.

Cases (n=252) were defined as children < 24 months of age with a conclusive diagnosis of acute lymphoid leukemia, ALL (n= 193), or acute myeloid leukemia, AML (n=59) confirmed by morphology, immunophenotype and standard cytogenetic-molecular methods.

Controls (n=423) were selected from children < 24 months of age with non-malignant diseases that were patients at the Brazilian National Health System oncologic centers where

cases were recruited, or patients of general hospitals in the same cities . These hospitals from which controls were recruited had the same catchment areas of those of cases. Controls included children with infectious and parasitic diseases (n=124, 29.4%); non-malignant hematological diseases (n=83, 19.6%); asthma and bronchitis (n=43, 10.2%); hemangioma (n=40, 9.4%); severe diarrhea (n=39, 9.2%); cardiovascular diseases (n= 25, 5.8%); and other non-malignant conditions (n=69, 16.4%).

Children with congenital syndromes, myelodysplasia, adoptive parents, or unknown biological mothers, were not eligible to be enrolled, and controls with a cancer diagnosis were excluded. Participation of invited cases and controls in the study was, respectively, 96 and 95% (Pombo-de-Oliveira and Koifman 2006).

Data Collection

The study was specifically designed to collect information on several environmental exposures potentially associated with leukemogenesis. Data were collected by in-person interviews with case and control mothers at the hospital. A standardized questionnaire was used for all participants, including information on the environmental exposures during pregnancy, including use of pesticides. Interviewers were health staff members without special knowledge of pesticide toxicology, and who were instructed to register all answers provided by the interviewed mothers. Pesticide exposure information was further analyzed and classified according to their toxicological characteristics. Child skin color was indicated by mothers and further analyzed as white and non-white. At each center, the same interviewers were responsible for both cases and controls. Participating mothers provided written informed consent for themselves and their child.

Pesticide exposure was evaluated based on the mother's report of any contact with pesticides (at least once) during the three months before pregnancy (periconceptual period),

throughout each pregnancy trimester, or during the three months after birth (breastfeeding). They were requested to inform about any contact with pesticides at home or in the workplace during each of these pregnancy times windows of exposure. Brand names of commercial products reported by the mothers were used to determine chemical content, and associations with pesticides were explored according to: the form of use, i.e., unintentional, domestic (household), or agricultural (maternal occupational exposure, or living in an agricultural area with pesticides use); and duration and regularity of contact (no use, \leq once a week, $>$ once a week).

Exposures to products that included pesticides from multiple chemical classes (for instance, pyrethroids, organophosphates and carbamates), or maternal exposure to both insecticides and herbicides were also assessed, and categorized as mixed exposures whenever reported.

Ethical Aspects

This investigation was approved by the Research Ethics Committee of the Brazilian National Cancer Institute (number No. 005/06) and by the Research Ethics Committee of the Oswaldo Cruz Foundation (FIOCRUZ), No. 32/10. Participating mothers provided written informed consent for themselves and their child.

Statistical Analysis

Unconditional logistic regression was performed to estimate associations between pesticide exposures and early leukemias based odds ratios and their 95% confidence intervals after adjustment for birth weight ($< 4,000$ g, $\geq 4,000$ g), maternal age at birth (< 35 years, ≥ 35 years), maternal schooling (≤ 8 years, > 8 years), oral contraceptive intake during pregnancy (no use, use during pregnancy), and child's skin color (white or non-white).

Sensitivity analysis was performed using different control subsets, specifically: a) after excluding controls with gastro-intestinal infections, parasitic diseases, dehydration, malnutrition or diarrhea; b) controls with respiratory illnesses, including tuberculosis, pneumonia, asthma, bronchitis, and bronchiolitis. Considering that both subsets are more prevalent among low income strata, such procedure aimed to evaluate possible confounding resulting from these controls inclusion. In addition, we estimated stratum-specific ORs according to child's skin color.

RESULTS

Origin of participants is presented at Supplemental Material, Table S1, showing a higher inclusion of children from Sao Paulo and Rio de Janeiro. Socio-demographic characteristics differed between cases and controls. Specifically, cases were more likely than controls to be white, to have mothers who were older at child's birth, more likely to have ≥ 8 years of education, and to have higher family income (Table 1). Complete information on model covariates was available for 85% of cases and 90% of controls. Agricultural pesticide exposures were reported by 22 (3.3%) mothers, including 7 (1.0%) who were agricultural workers.

Pesticide use at any time during the pregnancy was reported by 60.7% of AML, 36.4% of ALL, and 21.3% of control mothers of children at 0–11 months of age (Table 2). Adjusted odds ratios were 2.10 (95% CI: 1.14, 3.86) for ALL and 5.01 (95% CI: 1.97, 12.7) for AML.

For children diagnosed or enrolled at 12–23 months of age, 48.4%, 47.6% and 31.4% of AML, ALL, and controls were exposed to pesticides during pregnancy, respectively. Adjusted odds ratios were 1.88 (95% CI: 1.05, 5.23) for ALL and 1.98 (95% CI: 0.83, 4.74) for AML.

Among children diagnosed or enrolled at 0–11 months of age, information on periconceptual pesticide exposure was available for 99% of controls, 92% of ALL and 82% of AML. At age 12–23 months, they were, respectively, 95%, 63% and 71% (Table 2). ALL and AML were significantly associated with periconceptual exposures among children 0–11 months of age (adj. OR 2.40; 95% CI: 1.20, 4.81 and adj. OR 3.81; 95% CI: 1.34, 10.8, respectively). AML was significantly associated with periconceptual exposure in children age 12–23 months (adj. OR 2.48; 95% CI: 1.20, 5.11).

The odds of AML were increased with exposure during all time periods among children 0–11 months, with significant associations for pesticide exposure in the third trimester (adj. OR 3.70; 95% CI: 1.32, 10.4) and during breastfeeding (adj. OR 7.04; 95% CI: 2.47, 20.1). At age 12–23 months, they were, respectively, adj. OR 0.97; 95% CI: 0.35, 2.69 and adj. OR 1.20; 95% CI: 0.43, 3.34.

Sensitivity analysis according to the variables skin color (Supplemental Material, Table S2), diarrhea, parasitic diseases, dehydration or malnutrition (N = 98, Supplemental Material, Table S3), and respiratory diseases (Supplemental Material, Table S4) seems to support the presented results on the association between pesticides exposure and leukemia among very young children (Table 2).

Adjusted ORs for any exposure to pyrethroid pesticides during pregnancy were 1.80 (95% CI: 1.11, 2.90) for ALL and 3.39 (95% CI: 1.72, 16.78) for AML (Table 3). Adjusted ORs for use of pesticide formulations that included solvents were 1.79 (95% CI: 1.10, 2.92) for ALL and 3.45 (95% CI: 1.76, 6.74) for AML (Table 3).

The reported use of pesticide brands containing organophosphates was uncommon (1.7% of controls, 2.6% of AML and 6.8% of ALL mothers). The magnitude of association between such products and AML in very young children was adj. OR 5.50 (95% CI: 1.44, 21.03), (Table 3).

Adjusted ORs for pesticide exposure at home were 3.12 (95% CI: 1.61, 6.05) for AML, and 1.88 (95% CI: 1.20, 2.95) for ALL (Table 3). ORs for agricultural pesticide exposures were indeed higher, adj. OR 5.25; 95% CI: 1.83, 15.08 for ALL, and OR 7.56; 95% CI: 1.83, 31.23 for AML.

Compared with no reported pesticide exposure, an adjusted OR 2.88 (95% CI: 1.35, 6.17) for AML was estimated for the offspring of women reporting exposure \leq once a week, and OR 2.94 (95% CI: 1.17, 7.38) for exposure $>$ once a week (Table 3). For ALL, corresponding ORs were 2.19 (95% CI: 1.29, 3.69) and 2.06 (95% CI: 1.10, 3.86). Adjusted ORs for exposure to commercial products that included multiple pesticides were 5.22 (95% CI: 1.44, 19.0) for ALL, and 6.51 (95% CI: 1.25, 34.0) for AML.

A comprehensive analysis of all individual chemical components included in the reported brands revealed statistically significant associations between leukemia in very young children and maternal exposure to seven pyrethroids and unspecified solvents (Table 4). Estimates of associations with exposure to individual pyrethroids during pregnancy were imprecise due to small numbers of reported exposures. Permethrin, imiprothrin and esbiothrin exposures during pregnancy were positively associated either with ALL (adj. OR 2.47; 95% CI: 1.17, 5.25; adj. OR 2.61; 95% CI: 1.06, 6.93; adj. OR 3.03; 95% CI: 1.13, 8.09, respectively) or AML (adj. OR 7.28; 95% CI: 2.60, 20.38; adj. OR 3.41; 95% CI: 0.98, 11.90; adj. OR 3.19 95% CI: 0.77, 13.19) at age 0-11 months.

Among children age 11-23 months, esbiothrin exposures during pregnancy were positively associated with AML (adj. OR 3.71; 95% CI: 1.18, 11.62), (Table 4). Maternal exposure to Tetramethrin and D-allothrin was positively associated with AML in children 0-11 months (adj. OR 6.19; 95% CI: 2.07, 18.56 for both), while D-phenothrin was positively associated with AML in children 11-23 months of age (adj. OR 8.43; 95% CI: 1.59, 44.75), (Table 4).

DISCUSSION

Pesticides are complex mixtures that include components such as solvents, humidifying agents, emulsifiers and additives, in addition to active ingredients (Bolognesi 2003; Feron et al. 1998). Furthermore, the seasonal use of distinct formulas for specific purposes makes it difficult to present a qualitative evaluation of exposure to individual substances.

Our findings suggest that children whose mothers were exposed to pesticides 3 months before conception were at least twice as likely to be diagnosed with ALL in the first year of life compared to those who did not report such exposure. Adjusted ORs for AML in the first year of life ranged from 2.75 (95% CI: 0.96, 7.92) for any pesticide exposure in the first trimester of pregnancy, to 7.04 (95% CI: 2.47, 20.10) for exposure during breastfeeding.

Studies conducted in other countries have also reported positive associations between pesticide exposure and hematopoietic neoplasms in children, especially leukemias and lymphomas (Ma et al. 2002; Meinert et al. 2000; Menegaux et al. 2006; Rudant et al. 2007; Zahm and Ward 1998). A systematic review and meta-analysis of 15 studies of the association between residential exposure to pesticides during selected time windows (preconception, pregnancy and childhood) and childhood leukemia carried out between 1950-2009 (Turner et al. 2010), reported associations with pregnancy exposure to unspecified pesticides (OR 1.54; 95% CI: 1.13, 2.11), insecticides (OR 2.05; 95% CI: 1.80, 2.32) and herbicides (OR 1.61; 95% CI: 1.20, 2.16). Another meta-analysis of 31 studies of parental occupational exposure to pesticides and childhood leukemia (Wigle et al. 2009) reported associations with occupational exposure to insecticides (OR 2.72; 95% CI: 1.47, 5.04) and herbicides (OR 3.62; 95% CI: 1.28, 10.3) during pregnancy.

A French study also examined the association between pesticide exposure and infant leukemia (Rudant et al. 2007). According to use of any pesticide, the observed risk estimates were OR 2.3; 95% CI: 1.9, 2.8 for ALL, and OR 2.2; 95% CI: 1.4, 3.3 for AML. These authors also suggested that a domestic use of pesticides may play a role in the etiology of leukemia, and that prenatal exposure may be a window of fetal vulnerability.

Pesticide exposure during childhood may occur in many ways, either through contamination of their parents' work clothes or through household residues in water, air, soil and food (Araújo et al. 2000; Rudant et al. 2000). However, the short latency period for leukemias diagnosed during the first year of life suggests that intrauterine exposures may play a paramount role in this process.

In the Agricultural Health Study, a large prospective cohort study of approximately 49,000 pesticides applicators in the USA, an association between permethrin exposure and multiple myeloma was observed (Rusiecki et al. 2009). Compared to applicators who reported never using permethrin, the risk ratio for multiple myeloma among applicators in the highest tertile of lifetime exposure was RR 5.72 (95% CI: 2.76, 11.87). Another study of this cohort (Flower et al. 2003) reported a positive association between having a parent who applied pesticides and lymphomas diagnosed among children, age standardized incidence ratio, SIR 2.18; 95% CI: 1.13, 4.19, but not leukemias in children, SIR 0.91; 95% CI: 0.47, 1.95.

Moreover, the US Environmental Protection Agency (EPA) and the Canadian Pest Management Regulatory Agency (PMRA) have referred the occurrence of carcinogenicity following permethrin exposure in animal toxicity studies (Weichental et al. 2010). An insecticide containing imiprothrin and deltamethrin that is widely used in Egypt has been evaluated for immunotoxic effects in rats (Emara et al. 2007). The authors observed that animals exposed to both chemicals, compared to unexposed obnes, had altered levels of

splenic CD4+CD8- and CD4+CD8+ cells, and concluded that a repeated noncontinuous inhalation of imiothropin and deltamethrin causes several immunotoxic effects in other distal sites to the lungs. Other pyrethroids, such as allethrin, cyhalothrin, cypermethrin, deltamethrin and tetramethrin, have also been suggested to be involved in canine mammary carcinogenesis (Andrade et al. 2010).

This research has some limitations. The hospital-based case-control study design may introduce selection bias depending on the chosen comparison groups. We recruited controls with a variety of indications for hospitalization and enrolled controls from general hospitals in the same cities, though not necessarily the same hospitals, in which the cases were diagnosed. On the other hand, the similar origin of cases and controls could theoretically induce the introduction of overmatching in relation to agriculture pesticides exposure. The reports on pesticides exposure in the agricultural set (15 ALL, 4 AML and 7 controls, Table 3) accounted for 3.3% of all participants. Therefore, we think that the occurrence of overmatching on agricultural pesticide exposure distorting the presented conclusions of this investigation is improbable.

Pesticide exposures during the examined time windows were highly correlated, with statistically significant high Pearson's correlation coefficients, $r > 0.77$. Hence, associations with exposures during specific time of windows could not be accurately determined. Additionally, length of exposure was not evaluated in this study, so that associations according to maternal cumulative exposure to pesticides could not be estimated. Finally, sample size was limited, mainly to AML, thus resulting in imprecise estimates of association.

On the other hand, the study has some strengths, being relatively large given that the outcomes are rare. In addition, most previous studies have been based on populations from a limited number of countries, thus our study contributes for exploring the role of pesticides exposure during pregnancy and leukemias in children <2 years of age. Moreover, information

on the type of pesticide exposure, time periods of exposure, exposures to individual chemicals (mainly pyrethroids), and data on subgroups of leukemias (ALL, AML), can possibly contribute to enhance understanding of the role of maternal exposure to pesticides during pregnancy and leukemia in young children.

CONCLUSIONS

Future research will benefit from exploring the genetic and molecular mechanisms that characterize individual susceptibility to pesticide exposures in the development of leukemia in young children. However, the consistency of our findings with similar studies performed in different populations, supports recommendations for women of reproductive age to minimize their exposure to pesticides before and during pregnancy and breastfeeding.

REFERENCES

Alexander FE, Patheal SL, Biondi A, Brandalise S, Cabrera ME, Chan LC, et al. 2001. Transplacental chemical exposure and risk of infant leukemia with MLL gene fusion. *Cancer Research* 61:2542-2546.

Andrade FH, Figueiroa FC, Bersano PR, Bissacot DZ, Rocha NS. 2010. Malignant mammary tumor in female dogs: environmental contaminants. *Diagn Pathol*. 5:45.

Araujo AC, Nogueira DP, Augusto LG. 2000. Pesticide impact on health: a study of tomato cultivation. *Rev. de Saúde Pública* 34:309-313.

Barletta E, Gorini G, Vineis P, Miligi L, Davico L, Mugnai G, et al. 2004. Ras gene mutations in patients with acute myeloid leukaemia and exposure to chemical agents. *Carcinogenesis* 25(5):749-755.

Bassil KL, Vakil C, Sanborn N, Cole DC, Kaur JS, Kerr KJ. 2007. Cancer health effects of pesticides: systematic review. *Can Fam Physician* 53:1704-1711.

Bolognesi C. 2003. Genotoxicity of pesticides: a review of human biomonitoring studies. *Mutation Research* 543: 251-272.

Buffler PA, Kwan ML, Reynolds P, Urayama KY. 2005. Environmental and genetic risk factors for childhood leukemia: appraising the evidence. *Cancer Invest* 23:60-75.

Colt JS, Blair A. 1998. Parental exposures and risk of childhood cancer. *Environ Health Perspect* 106:909-925.

Emara AM, Draz EI. 2007. Immunotoxicological study of one of the most common over-the-counter pyrethroid insecticide products in Egypt. *Inhal Toxicol* 19:997-1009.

Emerenciano M, Agudelo Arias DP, Coser VM, de Brito GD, Macedo Silva ML, Pombo-de-Oliveira MS. 2006. Molecular cytogenetic findings of acute leukemia included in the

Brazilian Collaborative Study Group of Infant acute leukemia. *Pediatr Blood Cancer* 47: 549-554.

Faria NMX, Rosa JAR, Facchini LA. 2009. Intoxicações por agrotóxicos entre trabalhadores rurais de fruticultura, Bento Gonçalves, RS. *Rev. Saúde Pública* 43(2):335-344.

Falck GC, Hirvonen A, Scarpato R, Saarikoski ST, Migliore L, Norppa H. 1999. Micronuclei in blood lymphocytes and genetic polymorphism for GSTM1, GSTT1 and NAT2 in pesticide-exposed greenhouse workers. *Mutation Research* 441:225-237.

Feron VJ, Cassee FR, Groten JP. 1998. Toxicology of chemicals mixtures: international perspective. *Environ. Health Perspect* 106 Suppl 6:1281-1289.

Gabbianelli R, Nasuti C, Falcioni G, Cantalamessa F. 2004. Lymphocyte DNA damage in rats exposed to pyrethroids: effect of supplementation with Vitamins E and C. *Toxicology* 15:17-26.

Infante-Rivard C, Weichenthal S. 2007. Pesticides and childhood cancer: an update of Zahm and Ward's 1998 review. *J Toxicol Environ Health B Crit Rev* 10:81-99.

Kasat K, Go V, Pogo BG. 2002. Effects of pyrethroid insecticides and estrogen on WNT10B proto-oncogene. *Environ Int* 28:429-432.

Khan NI, Bradstock KF, Bendall LJ. 2007. Activation of Wnt/beta-catenin pathway mediates growth and survival in B-cell progenitor acute lymphoblastic leukaemia. *Br J Haematol* 138:338-348.

Kirikoshi H, Sekihara H, Katoh M. 2001. WNT10A and WNT6, clustered in human chromosome 2q35 region with head-to-tail manner, are strongly coexpressed in SW480 cells. *Biochem Biophys Res Commun* 283:798-805.

Lafiura KM, Bielawski DM, Posecion NC Jr, Ostrea E M Jr, Matherly LH Taub JW et al. 2007. Association between prenatal pesticide exposures and the generation of leukemia-associated t(8;21). *Pediatr Blood Cancer* 49:624–628.

Landrigan PJ, Claudio L, Markowitz SB, Berkowitz GS, Brenner BL, Romero H, Wetmur JG, Matte TD, Gore AC, Godbold JH, Wolff MS. 1999. Pesticides and inner-city children: exposures, risks, and prevention. *Environ Health Perspect*. 107(Suppl 3): 431–437.

Ma X, Buffler PA, Gunier RB, Dahl G, Smith MT, Renier K, et al. 2002. Critical windows of exposure to household pesticides and risk of acute leukemia. *Environ Health Perspect* 110:955-960.

Meyer C, Kowarz E, Hofmann J, Renneville A, Zuna J, Trka J, et al. 2009. New insights to the MLL recombinome of acute leukemias. *Leukemia* 23:1490-1499.

Meinert R, Schuz J, Kaletsch U, Kaatsch Michaelis J. 2000. Leukemia and non-Hodgking's lymphoma in childhood and exposure to pesticides: results of a register-based case-control Study in Germany. *Am J Epidemiol* 151:639-646.

Menegaux F, Baruchel A, Bertrand Y, Lescoeur B, Leveger G, Nelken B, et al. 2006. Household exposure to pesticides and risk of childhood acute leukaemia. *Occup Environ Med* 63:131-134.

Miller JR. 2001. The Wnts. *Genome Biol* 3:review3001.1-3001.15.

Mitrunen K, Jourenkova N, Kataja V, Eskelinen M, Kosma VM, Benhamou S, et al. 2001. Glutathione S-transferase M1, M3, P1, and T1 genetic polymorphisms and susceptibility to breast cancer. *Cancer Epidemiology Biomarkers Prevention* 10:229-236.

Monge P, Wesseling C, Guardado J, Lundberg I, Ahlbom A, Cantor KP et al. 2007. Parental occupational exposure to pesticides and the risk of childhood leukemia in Costa Rica. *Scand J Work Environ Health* 33:293-303.

Monge P, Partanen T, Wesseling C, Bravo V, Ruepert C, Burstyn I. 2005. Assessment of Pesticides Exposure in the Agricultural Population of Costa Rica. *Am. Occup Hyg* 49:375-384.

Perera FP. 1996. Molecular epidemiology: insights into cancer susceptibility, risk assessment, and prevention. *J Nat Cancer Inst* 88:496-509.

Perez-Saldivar ML, Ortega-Alvarez MC, Fajardo-Gutierrez A, Bernaldez-Rios R, Del Campo-Martinez ML, Medina-Sanson A, et al. 2008. Father's occupational exposure to carcinogenic agents and childhood acute leukemia: a new method to assess exposure (a case-control study). *BMC Cancer* 8:7.

Pombo-de-Oliveira MS, Koifman S, Brazilian Colaborative Study Group of Infant Leukemia. 2006. Infant acute leukemia and maternal exposures during pregnancy. *Cancer Epidemiol Biomarkers Prev* 15:2336-2341.

Raunio H, Husgafvel-Pursianen K, Antilla S, Hietanen E, Hirvonen NA, Pelkonen O. 1995. Diagnosis of polymorphisms in carcinogen-activating and inactivating enzymes and cancer susceptibility - a review. *Gene* 159:113-121.

Rudant J, Menegaux F, Leverger G, 2007. Household exposure to pesticides and risk of childhood hematopoietic malignancies: the SCALE study (SFCE). *Environ Health Perspect* 115:1787-1793.

Rudant J, Clavel J, Infant-Rivard C. 2010. Selection bias in case-control studies in household exposure to pesticides and childhood acute leukemia. *J Expo Sci Environ Epidemiol* 20:299-309.

Rusiecki JA, Patel R, Koutros S, Beane-Freeman L, Landgren O, Bonner MR, et al. 2009. Cancer incidence among pesticide applicators exposed to permethrin in the Agricultural Health Study. *Environ Health Perspect* 117:581-586.

Turner MC, Wigle DT, Krewski D. 2010. Residential pesticides and childhood leukemia: a systematic review and meta-analysis. *Environ Health Perspect* 118:33-41.

Wacholder S, McLaughlin JK, Silverman DT, Mandel JS. 1992. Selection of controls in case-control studies. I. Principles. *Am J Epidemiol* 135:1019-1028.

Weichenthal S, Moase C, Chan P. 2010. A review of pesticide exposure and cancer incidence in the Agricultural Health Study cohort. *Environ Health Perspect* 118:1117-1125.

Wigle DT, Turner MC, Krewski D. 2009. A systematic review and meta-analysis of childhood leukemia and parental occupational pesticide exposure. *Environ Health Perspect* 117:1505-1513.

Zahm SH, Ward MH. 1998. Pesticides and childhood cancer. *Environ Health Perspect* 106:893-908.

Table 1 - Distribution of selected socio-demographic variables, leukemia cases and controls, children < 2 yr., Brazil, 1999-2007

Characteristics		Cases (n=252) N (%)	Controls (n=423) N (%)	p-value
Sex				
	Male	130 (51.6)	226 (53.4)	0.643
	Female	122 (48.4)	197 (46.6)	
	Missing	0	0	
Birth weight				
	<4000g	234 (92.9)	393 (93.0)	0.470
	≥ 4000g	16 (6.3)	21 (5.0)	
	Missing	2 (0.8)	9 (2.0)	
Child's skin color				
	White	170 (67.5)	153 (36.2)	< 0.001
	Non-white	77 (30.5)	256 (60.5)	
	Missing	5 (2.0)	14 (3.3)	
Place of Birth				
	Northeast	52 (20.6)	101 (24.1)	0.552
	Midwest	18 (7.1)	31 (7.3)	
	Southeast	155 (61.5)	237 (56.0)	
	South	27 (10.7)	54 (12.6)	
	Missing	0	0	
Maternal Age (yr.) ^a				
	<18	8 (3.2)	60 (14.1)	< 0.001
	18-24	91 (36.1)	182 (42.9)	
	25-34	117 (46.4)	145 (34.2)	
	>35	36 (14.3)	36 (8.8)	
	Missing	0	0	
Maternal Education (yr.)				
	<8	81 (32.1)	206 (48.6)	< 0.001
	≥8	146 (57.9)	209 (49.4)	
	Missing	25 (10.0)	8 (2.0)	
Family Income ^b				
	≤350	74 (29.4)	148 (35.0)	< 0.001
	351-1750	102 (40.5)	218 (51.5)	
	1751-3500	27 (10.7)	11 (2.6)	
	≥3500	12 (5.0)	3 (0.7)	
	Missing	37 (14.4)	43 (10.2)	
Oral Contraceptive Use				
	No	223 (88.5)	404 (95.5)	< 0.001
	Yes	29 (11.5)	19 (4.5)	
	Missing	0	0	

Abbreviations: a-maternal age at birth; b- income in Brazilian currency, Real (R\$)

Table 2 – Maternal exposure to pesticides by time window of exposure, leukemia cases and controls, children < 2 yr., Brazil, 1999-2007

Pesticide Exposure	Controls (n=423), n (%)	ALL (n=193), n (%)	AML (n=59), n (%)	ALL		AML	
				Crude OR (95% CI)	Adj OR ^a (95% CI)	Crude OR (95% CI)	Adj OR ^a (95% CI)
Pesticide use							
0-11 months							
No	200 (78.7)	56 (63.6)	11 (39.3)	1.00	1.00	1.00	1.00
Yes	54 (21.3)	32 (36.4)	17 (60.7)	2.12 (1.25-3.59)	2.10 (1.14-3.86)	5.72 (2.53-12.94)	5.01 (1.97-12.75)
Missing	0	0	0				
12-23 months							
No	116 (68.6)	55 (52.4)	16 (51.6)	1.00	1.00	1.00	1.00
Yes	53 (31.4)	50 (47.6)	15 (48.4)	1.99 (1.20-3.29)	1.88 (1.05-5.23)	2.05 (0.95-4.46)	1.98 (0.83-4.74)
Missing	0	0	0				
Periconceptual ^b							
0-11 months							
No	220 (86.6)	63 (71.6)	15 (53.6)	1.00	1.00	1.00	1.00
Yes	32 (12.6)	18 (20.4)	8 (28.6)	1.96 (1.03-3.73)	2.40 (1.20-4.81)	3.67 (1.44-9.34)	3.81 (1.34-10.84)
Missing	2 (0.8)	7 (8.0)	5 (17.8)				
12-23 months							
No	125 (74.0)	64 (61.0)	15 (48.4)	1.00	1.00	1.00	1.00
Yes	36 (21.3)	32 (30.5)	7 (22.6)	1.62 (0.61-4.28)	1.34 (0.47-3.85)	1.94 (1.04-3.61)	2.48 (1.20-5.11)
Missing	8 (4.7)	9 (8.5)	9 (29.0)				

Table 2 (continued)

Pesticide Exposure	Controls (n=423), n (%)	ALL (n=193), n (%)	AML (n=59), n (%)	ALL		AML	
				Crude OR (95% CI)	Adj OR ^a (95% CI)	Crude OR (95% CI)	Adj OR ^a (95% CI)
1 st Trimester							
0-11 months							
No	219 (86.2)	63 (71.6)	16 (57.2)	1.00	1.00	1.00	1.00
Yes	32 (12.6)	18 (20.4)	7 (25.0)	1.96 (1.03-3.72)	1.86 (0.94-3.72)	2.99 (1.14-7.84)	2.75 (0.96-7.92)
Missing	3 (1.2)	7 (8.0)	5 (17.8)				
12-23 months							
No	127 (75.1)	65 (61.9)	20 (64.5)	1.00	1.00	1.00	1.00
Yes	35 (25.0)	31 (29.5)	8 (25.8)	1.73 (0.98-3.05)	1.87 (0.99-3.56)	1.45 (0.59-3.57)	1.28 (0.47-3.53)
Missing	7 (4.1)	9 (8.5)	3 (9.7)				
2 nd Trimester							
0-11 months							
No	219 (86.2)	64 (72.7)	16 (57.2)	1.00	1.00	1.00	1.00
Yes	32 (12.6)	17 (19.3)	7 (25.0)	1.70 (0.89-3.25)	1.75 (0.87-3.55)	2.80 (1.08-7.32)	2.27 (0.79-6.47)
Missing	3 (1.2)	7 (8.0)	5 (17.8)				
12-23 months							
No	129 (76.2)	68 (64.8)	20 (64.5)	1.00	1.00	1.00	1.00
Yes	33 (19.5)	28 (26.7)	8 (25.8)	1.61 (0.90-2.88)	1.76 (0.91-3.39)	1.56 (0.63-3.86)	1.48 (0.54-4.04)
Missing	7 (4.1)	9 (8.5)	3 (9.7)				
3 rd Trimester							
0-11 months							
No	218 (85.8)	65 (73.8)	15 (53.6)	1.00	1.00	1.00	1.00
Yes	33 (13.0)	16 (18.2)	8 (28.6)	1.63 (0.84-3.14)	1.88 (0.93-3.79)	3.52 (1.39-8.96)	3.70 (1.32-10.38)
Missing	3 (1.2)	7 (8.0)	5 (17.8)				

Table 2 (continued)

Pesticide Exposure	Controls (n=423), n (%)	ALL (n=193), n (%)	AML (n=59), n (%)	ALL		AML	
				Crude OR (95% CI)	Adj OR ^a (95% CI)	Crude OR (95% CI)	Adj OR ^a (95% CI)
3 rd Trimester							
12-23 months							
No	123 (72.8)	68 (64.8)	20 (64.5)	1.00	1.00	1.00	1.00
Yes	39 (23.1)	28 (26.7)	8 (25.8)	1.30 (0.74-2.29)	1.26 (0.66-2.40)	1.26 (0.52-3.09)	0.97 (0.35-2.69)
Missing	7 (4.1)	9 (8.5)	3 (9.7)				
Breastfeeding ^c							
0-11 months							
No	229 (90.2)	69 (78.4)	14 (50.0)	1.00	1.00	1.00	1.00
Yes	22 (8.6)	12 (13.6)	9 (32.2)	1.81 (0.85-3.84)	2.05 (0.92-4.58)	6.69 (2.60-17.21)	7.04 (2.47-20.10)
Missing	3 (1.2)	7 (8.0)	5 (17.8)				
12-23 months							
No	128 (75.7)	68 (64.8)	21 (67.7)	1.00	1.00	1.00	1.00
Yes	33 (19.6)	28 (26.7)	7 (22.6)	1.60 (0.89-2.86)	1.53 (0.80-2.95)	1.29 (0.51-3.30)	1.20 (0.43-3.34)
Missing	8 (4.7)	9 (8.5)	3 (9.7)				

Abbreviations:
^a adjusted odds ratio (adj. OR) by use of oral contraceptives during pregnancy, maternal age and education, infant birth weight and child's skin color;
^b 3 months before pregnancy;
^c 3 months after delivery

Table 3 - Maternal use of pesticides according to chemical class, frequency and type of use along pregnancy, leukemia cases and controls, children < 2 yr., Brazil, 1999-2007

Use of pesticides along pregnancy	Controls (N=423) N (%)	ALL (N=193), n (%)	AML (N=59), n (%)	ALL vs controls Crude OR (95% CI)	ALL vs Controls adj OR (95% C.I.) ^a	AML vs controls Crude OR (95% CI)	AML vs Controls adj OR ^a (95% CI)
Chemical Groups							
No use	316 (74.7)	111 (57.5)	27 (45.7)	1.00	1.00	1.00	1.00
Pyrethroid	89 (21.0)	63 (32.6)	25 (42.4)	2.02 (1.34-3.02)	1.80 (1.11-2.90)	3.29 (1.73-6.20)	3.39 (1.72-16.78)
Organophosphates	7 (1.7)	5 (2.6)	4 (6.8)	2.03 (0.63-6.54)	1.06 (0.26-4.32)	6.69 (1.84-24.29)	5.50 (1.44-21.03)
Other Pesticides ^b	17 (2.6)	14 (7.3)	3 (5.1)	2.66 (1.30-5.43)	2.96 (1.28-6.84)	2.66 (0.84-8.44)	1.77 (0.35-8.80)
Type of use							
No use	316 (74.7)	111 (57.5)	27 (48.2)	1.00	1.00	1.00	1.00
Household	89 (21.0)	62 (32.1)	25 (44.6)	1.98 (1.34-2.93)	1.88 (1.20-2.95)	3.29 (1.82-5.95)	3.12 (1.61-6.05)
Agriculture	7 (1.7)	15 (7.8)	4 (7.1)	6.10 (2.42-15.35)	5.25 (1.83-15.08)	6.69 (1.84-24.29)	7.56 (1.83-31.23)
Frequency							
None	315 (79.3)	111 (64.9)	27 (55.1)	1.00	1.00	1.00	1.00
Up to once/week	45 (11.3)	35 (20.5)	13 (26.5)	2.23 (1.38-3.59)	2.19 (1.29-3.69)	3.20 (1.58-6.48)	2.88 (1.35-6.17)
> once/week	37 (8.7)	25 (13.0)	9 (15.3)	1.82 (1.02-3.25)	2.06 (1.10-3.86)	2.58 (1.09-6.09)	2.94 (1.17-7.38)
Brands with mixed chemicals^c							
No	414 (97.9)	183 (94.8)	55 (93.2)	1.00	1.00	1.00	1.00
Yes	9 (2.1)	10 (5.2)	4 (6.8)	2.51 (1.01-6.29)	5.22 (1.44-18.97)	3.35 (0.73-12.35)	6.51 (1.25-33.99)
Distinct pest classes							
No	420 (99.3)	186 (96.4)	58 (98.3)	1.00	1.00	1.00	1.00
Yes	3 (0.7)	7 (3.6)	1 (1.7)	5.27 (1.35-20.6)	18.34 (2.01-167.38)	3.35 (1.00-11.23)	8.03 (0.37-174.68)

a- Adjusted odds ratio (adj. OR) by use of oral contraceptives during pregnancy, maternal age, maternal education, birth weight and child's skin color b- Organochlorines, cumarines and others.

c- Exposure to more than one chemical in the same product or distinct products.

Table 4 – Maternal exposure to specific pesticides chemical components along pregnancy, leukemia cases and controls, children < 2 yr., Brazil, 1999-2007

Chemical compounds	Controls	ALL	AML	ALL		AML	
	(N=423) n (%)	(N=193) n (%)	(N=59) n (%)	Crude OR (95% CI)	Adj OR ^a (95% C.I.)	Crude OR (95% CI)	Adj OR ^a (95% CI)
Prallethrin							
0-11 months							
No	234 (55.3)	78 (40.4)	23 (43.4)	1.00	1.00	1.00	1.00
Yes	3 (0.7)	1 (0.5)	2 (3.8)	1.00 (0.10-9.75)	1.52 (0.15-15.32)	6.78 (1.08-42.70)	8.06 (1.17-55.65)
12-23 months							
No	155 (36.6)	94 (48.7)	29 (54.7)	1.00	1.00	1.00	1.00
Yes	3 (0.7)	3 (1.5)	0 (0.0)	1.65 (0.33-8.34)	1.16 (0.15-9.12)	-	-
Permethrin							
0-11 months							
No	211 (49.9)	63 (32.6)	15 (28.3)	1.00	1.00	1.00	1.00
Yes	26 (6.1)	16 (8.3)	10 (18.9)	2.06 (1.04-4.08)	2.47 (1.17-5.25)	5.41 (2.20-13.28)	7.28 (2.60-20.38)
12-23 months							
No	128 (30.3)	75 (38.8)	23 (43.4)	1.00	1.00	1.00	1.00
Yes	30 (7.1)	22 (11.4)	6 (11.3)	1.25 (0.67-2.33)	1.47 (0.71-3.08)	1.11 (0.42-2.97)	1.32 (0.43-4.02)
Imiprothrin							
0-11 months							
No	223 (52.7)	67 (34.7)	20 (37.7)	1.00	1.00	1.00	1.00
Yes	14 (3.3)	12 (6.2)	5 (9.4)	2.85 (1.26-6.46)	2.61 (1.06-6.93)	3.98 (1.30-12.91)	3.41 (0.98-11.90)
12-23 months							
No	140 (33.1)	81 (42.0)	22 (41.5)	1.00	1.00	1.00	1.00
Yes	18 (4.3)	16 (8.3)	7 (13.2)	1.54 (0.74-3.18)	1.38 (0.59-3.23)	2.48 (0.93-6.61)	2.85 (0.94-8.62)

Table 4 (continued)

Chemical compounds	Controls (N=423) n (%)	ALL (N=193) n (%)	AML (N=59) n (%)	ALL		AML		
				Crude OR (95% CI)	Adj OR ^a (95% C.I.)	Crude OR (95% CI)	Adj OR ^a (95% CI)	
Esbiothrin								
0-11 months								
No	227 (53.7)	68 (35.2)	21 (39.6)	1.00	1.00	1.00	1.00	
Yes	10 (2.4)	11 (5.7)	4 (7.5)	3.67 (1.50-9.02)	3.03 (1.13-8.09)	4.32 (1.25-14.98)	3.19 (0.77-13.19)	
12-23 months								
No	144 (34.0)	83 (43.0)	22 (41.5)	1.00	1.00	1.00	1.00	
Yes	14 (3.3)	14 (7.3)	7 (13.2)	1.74 (0.79-3.82)	1.66 (0.67-4.13)	3.27 (1.19-9.00)	3.71 (1.18-11.62)	
Tetramethrin								
0-11 months								
No	214 (50.6)	68 (35.2)	17 (32.1)	1.00	1.00	1.00	1.00	
Yes	23 (5.4)	11 (5.7)	8 (15.1)	1.51 (0.70-3.25)	1.56 (0.65-3.72)	4.38 (1.70-11.25)	6.19 (2.07-18.56)	
12-23 months								
No	134 (31.7)	78 (40.4)	27 (50.9)	1.00	1.00	1.00	1.00	
Yes	24 (5.7)	17 (8.8)	2 (3.8)	1.22 (0.66-2.40)	1.35 (0.59-3.07)	0.41 (0.09-1.85)	0.47 (0.09-2.48)	
D-Phenothrin								
0-11 months								
No	234 (55.3)	72 (37.3)	24 (45.3)	1.00	1.00	1.00	1.00	
Yes	3 (0.7)	7 (3.6)	1 (1.9)	7.58 (1.91-30.08)	4.16 (0.85-20.29)	3.25 (0.33-32.48)	1.64 (0.16-19.68)	
12-23 months								
No	155 (36.6)	93 (48.2)	25 (47.2)	1.00	1.00	1.00	1.00	
Yes	3 (0.7)	4 (2.1)	4 (7.5)	2.22 (0.49-10.15)	0.69 (0.10-4.88)	8.27 (1.75-39.16)	8.43 (1.59-44.75)	

Table 4 (continued)

Chemical compounds	Controls	ALL	AML	ALL		AML	
	(N=423) n (%)	(N=193) n (%)	(N=59) n (%)	Crude OR (95% CI)	Adj OR ^a (95% C.I.)	Crude OR (95% CI)	Adj OR ^a (95% CI)
D-Allethrin							
0-11 months							
No	213 (50.3)	68 (35.2)	17 (32.1)	1.00	1.00	1.00	1.00
Yes	24 (5.7)	11 (5.7)	8 (15.1)	1.43 (0.67-3.08)	1.56 (0.65-3.72)	4.18 (1.63-10.70)	6.19 (2.07-18.56)
12-23 months							
No	132 (31.2)	78 (40.4)	27 (50.9)	1.00	1.00	1.00	1.00
Yes	26 (6.1)	19 (9.8)	2 (3.8)	1.24 (0.64-2.38)	1.54 (0.70-3.39)	0.38 (0.08-1.68)	0.46 (0.09-2.40)
Table 4 continued B							
Solvents							
0-11 months							
No	205 (48.5)	59 (30.1)	13 (24.5)	1.00	1.00	1.00	1.00
Yes	32 (7.5)	20 (10.4)	12 (22.6)	2.17 (1.16-4.07)	2.17 (1.06-4.43)	5.91 (2.48-14.10)	6.70 (2.50-17.97)
12-23 months							
No	122 (28.8)	70 (36.3)	20 (37.7)	1.00	1.00	1.00	1.00
Yes	36 (8.5)	27 (14.0)	9 (17.0)	1.31 (0.73-2.33)	1.32 (0.66-2.63)	1.52 (0.64-3.64)	1.82 (0.68-4.84)

Abbreviations:
^a adjusted odds ratio (adj. OR) by use of oral contraceptives during pregnancy, maternal age, maternal education, birth weight and child’s skin color.